

$XYZ_w=95.04, 100.0, 108.89$

$A_2 = 2.5(a_2 - a_{2w})Y$

$B_2 = 2.5B_1(b_2 - b_{2w})Y$

$a_2 = a_{2w}[(x - x_c)/y]$

$b_2 = b_{2w}[z/y]$

$a_{2w} = 1, b_{2w} = -0.4$

$x_c = 0.110, B_1 = 0.800$


$C_{AB} = [A_2^2 + B_2^2]^{1/2}$

6 Oswald colours (o)

of maximum (m) C_{AB} in chromatic value diagram (A_2, B_2)

Illumin. D65, $Y_w=100, Y_N=50$

Table with 10 columns: Name, Range, X₁, Y₁, Z₁, X₂, Y₂, Z₂, X₃, Y₃, Z₃. Rows include 77G_d, 81C_d, 69R_d, 72M_d, 53B_d.



$XYZ_w=96.42, 100.0, 82.49$

$A_2 = 2.5(a_2 - a_{2w})Y$

$B_2 = 2.5B_1(b_2 - b_{2w})Y$

$a_2 = a_{2w}[(x - x_c)/y]$

$b_2 = b_{2w}[z/y]$

$a_{2w} = 1, b_{2w} = -0.4$

$x_c = 0.110, B_1 = 1.000$


$C_{AB} = [A_2^2 + B_2^2]^{1/2}$

6 Oswald colours (o)

of maximum (m) C_{AB} in chromatic value diagram (A_2, B_2)

Illumin. D50, $Y_w=100, Y_N=50$

Table with 10 columns: Name, Range, X₁, Y₁, Z₁, X₂, Y₂, Z₂, X₃, Y₃, Z₃. Rows include 77G_d, 80C_d, 70R_d, 73M_d, 53B_d.



$XYZ_w=100.93, 100.0, 64.68$

$A_2 = 2.5(a_2 - a_{2w})Y$

$B_2 = 2.5B_1(b_2 - b_{2w})Y$

$a_2 = a_{2w}[(x - x_c)/y]$

$b_2 = b_{2w}[z/y]$

$a_{2w} = 1, b_{2w} = -0.4$

$x_c = 0.110, B_1 = 1.300$


$C_{AB} = [A_2^2 + B_2^2]^{1/2}$

6 Oswald colours (o)

of maximum (m) C_{AB} in chromatic value diagram (A_2, B_2)

Illumin. F40, $Y_w=100, Y_N=50$

Table with 10 columns: Name, Range, X₁, Y₁, Z₁, X₂, Y₂, Z₂, X₃, Y₃, Z₃. Rows include 77G_d, 79C_d, P40, 71R_d, 73M_d, 52B_d.



$XYZ_w=109.84, 99.99, 35.58$

$A_2 = 2.5(a_2 - a_{2w})Y$

$B_2 = 2.5B_1(b_2 - b_{2w})Y$

$a_2 = a_{2w}[(x - x_c)/y]$

$b_2 = b_{2w}[z/y]$

$a_{2w} = 1, b_{2w} = -0.4$

$x_c = 0.110, B_1 = 2.500$


$C_{AB} = [A_2^2 + B_2^2]^{1/2}$

6 Oswald colours (o)

of maximum (m) C_{AB} in chromatic value diagram (A_2, B_2)

Illumin. A00, $Y_w=100, Y_N=50$

Table with 10 columns: Name, Range, X₁, Y₁, Z₁, X₂, Y₂, Z₂, X₃, Y₃, Z₃. Rows include 77G_d, 79C_d, A00, 72R_d, 74M_d, 52B_d.



$XYZ_w=100.0, 100.0, 100.0$

$A_2 = 2.5(a_2 - a_{2w})Y$

$B_2 = 2.5B_1(b_2 - b_{2w})Y$

$a_2 = a_{2w}[(x - x_c)/y]$

$b_2 = b_{2w}[z/y]$

$a_{2w} = 1, b_{2w} = -0.4$

$x_c = 0.110, B_1 = 0.900$


$C_{AB} = [A_2^2 + B_2^2]^{1/2}$

6 Oswald colours (o)

of maximum (m) C_{AB} in chromatic value diagram (A_2, B_2)

Illumin. E00, $Y_w=100, Y_N=50$

Table with 10 columns: Name, Range, X₁, Y₁, Z₁, X₂, Y₂, Z₂, X₃, Y₃, Z₃. Rows include 77G_d, 80C_d, E00, 70R_d, 73M_d, 53B_d.



$XYZ_w=98.07, 100.0, 118.22$

$A_2 = 2.5(a_2 - a_{2w})Y$

$B_2 = 2.5B_1(b_2 - b_{2w})Y$

$a_2 = a_{2w}[(x - x_c)/y]$

$b_2 = b_{2w}[z/y]$

$a_{2w} = 1, b_{2w} = -0.4$

$x_c = 0.110, B_1 = 0.700$


$C_{AB} = [A_2^2 + B_2^2]^{1/2}$

6 Oswald colours (o)

of maximum (m) C_{AB} in chromatic value diagram (A_2, B_2)

Illumin. C00, $Y_w=100, Y_N=50$

Table with 10 columns: Name, Range, X₁, Y₁, Z₁, X₂, Y₂, Z₂, X₃, Y₃, Z₃. Rows include 77G_d, 80C_d, C00, 70R_d, 73M_d, 53B_d.



$XYZ_w=102.06, 100.0, 81.06$

$A_2 = 2.5(a_2 - a_{2w})Y$

$B_2 = 2.5B_1(b_2 - b_{2w})Y$

$a_2 = a_{2w}[(x - x_c)/y]$

$b_2 = b_{2w}[z/y]$

$a_{2w} = 1, b_{2w} = -0.4$

$x_c = 0.110, B_1 = 1.000$


$C_{AB} = [A_2^2 + B_2^2]^{1/2}$

6 Oswald colours (o)

of maximum (m) C_{AB} in chromatic value diagram (A_2, B_2)

Illumin. P00, $Y_w=100, Y_N=50$

Table with 10 columns: Name, Range, X₁, Y₁, Z₁, X₂, Y₂, Z₂, X₃, Y₃, Z₃. Rows include 77G_d, 80C_d, P00, 70R_d, 73M_d, 53B_d.



$XYZ_w=97.93, 100.0, 118.95$

$A_2 = 2.5(a_2 - a_{2w})Y$

$B_2 = 2.5B_1(b_2 - b_{2w})Y$

$a_2 = a_{2w}[(x - x_c)/y]$

$b_2 = b_{2w}[z/y]$

$a_{2w} = 1, b_{2w} = -0.4$

$x_c = 0.110, B_1 = 0.700$


$C_{AB} = [A_2^2 + B_2^2]^{1/2}$

6 Oswald colours (o)

of maximum (m) C_{AB} in chromatic value diagram (A_2, B_2)

Illumin. Q00, $Y_w=100, Y_N=50$

Table with 10 columns: Name, Range, X₁, Y₁, Z₁, X₂, Y₂, Z₂, X₃, Y₃, Z₃. Rows include 78G_d, 81C_d, Q00, 70R_d, 72M_d, 53B_d.



BEH0-1R

BEH0-1A