

$XYZ_W = 108.04, 100.0, 39.55$

$A_2 = 2.5 (a_2 - a_{2,n}) Y$

$B_2 = 2.5 B_c (b_2 - b_{2,n}) Y$

$a_2 = a_{20} [(x - x_c) / y]$

$b_2 = b_{20} [z / y]$

$a_{20} = 1, b_{20} = -0.4$

$x_c = 0.110, B_c = 2.500$

$C_{AB2} = [A_2^2 + B_2^2]^{1/2}$

6 Ostwald colours (o)


of maximum (m) C_{AB} in

linear colour space ($C_{AB,2n} Y$)

Illumin. P30, $Y_W = 100, Y_N = 50$

Name	Range	X_d	Y_d	Z_d	x_d	y_d	λ_d	λ_c
R_d	578_775	93.17	71.4	19.83	0.5052	0.3871	604	498
Y_d	503_775	105.07	98.0	21.09	0.4687	0.4371	580	473
G_d	503_578	66.03	76.7	21.07	0.4031	0.4682	546	546c
C_d	380_578	69.05	78.74	39.54	0.3685	0.4203	498	604
B_d	380_503	57.14	52.14	38.28	0.3872	0.3533	473	580
M_d	578_503	96.19	73.44	38.3	0.4625	0.3531	546c	546
W_d	380_775	108.04	100.0	39.55	0.4363	0.4038	100%	
N_d	380_775	54.02	50.0	19.77	0.4363	0.4038	50%	
Z_d	380_775	19.44	18.0	7.11	0.4363	0.4038	18%	

$L^* = 60 \log[f(Y_{an})]$



$f(Y_{an}) = \pm [1 + 10 |Y_{an}|^n]$

n increases to 1 for:

1. decreasing of the contrast C
2. adjacent compared to separate colours.

Parameter:

Y & Name

Illuminant P30

$Y_W = 100, Y_N = 50$