

$XYZ_W = 100.0, 100.0, 100.0$

$A_2 = 2.5 (a_2 - a_{2,n}) Y$

$B_2 = 2.5 B_c (b_2 - b_{2,n}) Y$

$a_2 = a_{20} [(x - x_c) / y]$

$b_2 = b_{20} [z / y]$

$a_{20} = 1, b_{20} = -0.4$

$x_c = 0.110, B_c = 0.900$

$C_{AB2} = [A_2^2 + B_2^2]^{1/2}$

6 Ostwald colours (o)


of maximum (m) C_{AB} in

linear colour space ($C_{AB,2}, Y$)

Illumin. E00, $Y_W = 100, Y_N = 0$

Name	Range	X_d	Y_d	Z_d	x_d	y_d	λ_d	λ_c
R_d	570_775	65.52	40.49	0.24	0.6165	0.381	598	489
Y_d	494_775	83.39	94.82	6.19	0.4522	0.5141	573	463
G_d	494_570	18.07	54.52	6.14	0.2295	0.6924	536	536c
C_d	380_570	34.67	59.7	99.95	0.1784	0.3072	489	598
B_d	380_494	16.8	5.37	94.0	0.1446	0.0462	463	573
M_d	570_494	82.12	45.67	94.05	0.3701	0.2058	536c	536
W_d	380_775	100.0	100.0	100.0	0.3333	0.3333	100%	
N_d	380_775	0.1	0.1	0.1	0.3332	0.3332	0%	
Z_d	380_775	18.0	18.0	18.0	0.3333	0.3333	18%	

$L^* = 60 \log[f(Y_{an})]$



$f(Y_{an}) = \pm [1 + 10 |Y_{an}|^n]$

n increases to 1 for:

1. decreasing of the contrast C
2. adjacent compared to separate colours.

Parameter:

Y & Name

Illuminant E00

$Y_W = 100, Y_N = 0$