

$XYZ_w = 102.06, 100.0, 81.06$

$A_2 = 2,5 (a_2 - a_{2,n}) Y$

$B_2 = 2,5 B_c (b_2 - b_{2,n}) Y$

$a_2 = a_{20} [(x - x_c) / y]$

$b_2 = b_{20} [z / y]$

$a_{20} = 1, b_{20} = -0,4$

$x_c = 0,110, B_c = 1,000$

$C_{AB2} = [A_2^2 + B_2^2]^{1/2}$

6 Ostwald colours (o)


of maximum (m) C_{AB} in

linear colour space ($C_{AB,2r} Y$)

Illumin. P00, $Y_w = 100, Y_n = 25$

Name	Range	X_d	Y_d	Z_d	x_d	y_d	λ_d	λ_c
R _d	572_775	77.3	55.51	20.38	0.5046	0.3623	600	491
Y _d	496_775	92.09	95.88	23.27	0.4359	0.4538	575	467
G _d	496_572	40.4	65.47	23.24	0.3129	0.507	541	541c
C _d	380_572	50.4	69.61	81.04	0.2506	0.3462	491	600
B _d	380_496	35.61	29.24	78.15	0.249	0.2044	467	575
M _d	572_496	87.3	59.65	78.18	0.3877	0.2649	541c	541
W _d	380_775	102.06	100.0	81.06	0.3604	0.3531	100%	
N _d	380_775	25.51	25.0	20.26	0.3604	0.3531	25%	
Z _d	380_775	18.37	18.0	14.59	0.3604	0.3531	18%	

$L^* = 60 \log[f(Y_{an})]$



$f(Y_{an}) = \pm [1 + 10 |Y_{an}|^n]$

- n increases to 1 for:
1. decreasing of the contrast C
 2. adjacent compared to separate colours.

Parameter:
 Y & Name
 Illuminant P00
 $Y_w = 100, Y_n = 25$