

$XYZ_{W,10}=101.75, 100.0, 64.44$

$A_{2,10} = 2,5 (a_{2,10} - a_{2,n,10}) Y_{10}$

$B_{2,10} = 2,5 B_c (b_{2,10} - b_{2,n,10}) Y_{10}$

$a_{2,10} = a_{20} [(x_{10} - x_c) / y_{10}]$

$b_{2,10} = b_{20} [z_{10} / y_{10}]$

$a_{20} = 1, b_{20} = -0,4$

$x_c = 0,110, B_c = 1,300$

$C_{AB,2,10} = [A_{2,10}^2 + B_{2,10}^2]^{1/2}$

6 Ostwald colours (o)


of maximum (m) $C_{AB,10}$ in

linear colour space ($C_{AB,2,10}, Y_{10}$)

Illumin. P40, $Y_{W,10}=100, Y_{N,10}=0$

Name	Range	$X_{d,10}$	$Y_{d,10}$	$Z_{d,10}$	$x_{d,10}$	$y_{d,10}$	λ_d	λ_c
R_d	569_775	71.44	43.5	0.12	0.6208	0.378	597	487
Y_d	492_775	91.3	93.44	3.86	0.484	0.4954	572	465
G_d	492_569	20.05	50.14	3.86	0.2708	0.677	535	535c
C_d	380_569	30.5	56.69	64.44	0.2011	0.3738	487	597
B_d	380_492	10.65	6.75	60.71	0.1363	0.0864	465	572
M_d	569_492	81.89	50.05	60.71	0.425	0.2598	535c	535
W_d	380_775	101.75	100.0	64.44	0.3822	0.3756	100%	
N_d	380_775	0.1	0.1	0.06	0.382	0.3755	0%	
Z_d	380_775	18.31	18.0	11.6	0.3822	0.3756	18%	

$L^*_{10} = 60 \log[f(Y_{10,an})]$ $Y_{10,an} = [Y_{10} - 50] / 50$



$f(Y_{10,an}) = \pm [1 + 10 |Y_{10,an}|^n]$

n increases to 1 for:

1. decreasing of the contrast C
2. adjacent compared to separate colours.

Parameter:
 Y_{10} & Name
 Illuminant P40
 $Y_{W,10}=100, Y_{N,10}=0$