


$XYZ_{W,10} = 97.65, 100.0, 118.42$
 $A_{2,10} = 2,5 (a_{2,10} - a_{2,n,10}) Y_{10}$
 $B_{2,10} = 2,5 B_c (b_{2,10} - b_{2,n,10}) Y_{10}$
 $a_{2,10} = a_{20} [(x_{10} - x_c) / y_{10}]$
 $b_{2,10} = b_{20} [z_{10} / y_{10}]$
 $a_{20} = 1, b_{20} = -0,4$
 $x_c = 0,110, B_c = 0,700$
 $C_{AB,2,10} = [A_{2,10}^2 + B_{2,10}^2]^{1/2}$

6 Ostwald colours (o)
 of maximum (m) $C_{AB,10}$ in
 linear colour space $(C_{AB,2,10} Y_{10})_{B_d}$

Illumin. Q00, $Y_{W,10} = 100, Y_{N,10} = 0$

Name	Range	$X_{d,10}$	$Y_{d,10}$	$Z_{d,10}$	$x_{d,10}$	$y_{d,10}$	λ_d	λ_c
R_d	561_775	60.93	39.91	0.23	0.6028	0.3948	593	481
Y_d	486_775	77.48	90.9	6.21	0.4437	0.5206	566	459
G_d	486_561	16.74	51.18	6.21	0.2258	0.6903	530	530c
C_d	380_561	36.9	60.28	118.42	0.1711	0.2796	481	593
B_d	380_486	20.35	9.29	112.44	0.1432	0.0654	459	566
M_d	561_486	81.1	49.01	112.44	0.3343	0.202	530c	530
W_d	380_775	97.65	100.0	118.42	0.3089	0.3163	100%	
N_d	380_775	0.09	0.1	0.11	0.3088	0.3162	0%	
Z_d	380_775	17.57	18.0	21.31	0.3089	0.3163	18%	



$f(Y_{10,an}) = \pm [1 + 10 |Y_{10,an}|^n]$
n increases to 1 for:
 1. decreasing of the contrast C
 2. adjacent compared to separate colours.

Parameter:
 Y_{10} & Name
Illuminant Q00
 $Y_{W,10} = 100, Y_{N,10} = 0$