

$XYZ_{W,10} = 97.65, 100.0, 118.42$   $L^*_{10} = 60 \log[f(Y_{10,an})]$   $Y_{10,an} = [Y_{10} - 50] / 50$

$A_{2,10} = 2,5 (a_{2,10} - a_{2,n,10}) Y_{10}$

$B_{2,10} = 2,5 B_c (b_{2,10} - b_{2,n,10}) Y_{10}$

$a_{2,10} = a_{20} [(x_{10} - x_c) / y_{10}]$

$b_{2,10} = b_{20} [z_{10} / y_{10}]$

$a_{20} = 1, b_{20} = -0,4$

$x_c = 0,110, B_c = 0,700$

$C_{AB,2,10} = [A_{2,10}^2 + B_{2,10}^2]^{1/2}$


6 Ostwald colours (o)

of maximum (m)  $C_{AB,10}$  in

linear colour space ( $C_{AB,2,10} Y_{10}$ )

Illumin. Q00,  $Y_{W,10} = 100, Y_{N,10} = 4$

Name	Range	$X_{d,10}$	$Y_{d,10}$	$Z_{d,10}$	$x_{d,10}$	$y_{d,10}$	$\lambda_d$	$\lambda_c$
$R_d$	561_775	62.37	42.26	4.85	0.5696	0.3859	593	481
$Y_d$	486_775	78.27	91.26	10.59	0.4345	0.5066	566	459
$G_d$	486_561	19.9	53.09	10.59	0.2381	0.6351	530	530c
$C_d$	380_561	39.28	61.84	118.42	0.1789	0.2816	481	593
$B_d$	380_486	23.37	12.84	112.68	0.157	0.0862	459	566
$M_d$	561_486	81.75	51.0	112.68	0.333	0.2078	530c	530
$W_d$	380_775	97.65	100.0	118.42	0.3089	0.3163	100%	
$N_d$	380_775	3.9	4.0	4.73	0.3089	0.3163	4%	
$Z_d$	380_775	17.57	18.0	21.31	0.3089	0.3163	18%	



$f(Y_{10,an}) = \pm [1 + 10 |Y_{10,an}|^n]$   
 n increases to 1 for:  
 1. decreasing of the contrast C  
 2. adjacent compared to separate colours.

Parameter:  
 $Y_{10}$  & Name  
 Illuminant Q00  
 $Y_{W,10} = 100, Y_{N,10} = 4$