

$XYZ_{W,10} = 102.37, 99.99, 81.25$

$A_{2,10} = 2.5 (a_{2,10} - a_{2,n,10}) Y_{10}$

$B_{2,10} = 2.5 B_c (b_{2,10} - b_{2,n,10}) Y_{10}$

$a_{2,10} = a_{20} [(x_{10} - x_c) / y_{10}]$

$b_{2,10} = b_{20} [z_{10} / y_{10}]$

$a_{20} = 1, b_{20} = -0.4$

$x_c = 0.110, B_c = 1.000$

$C_{AB,2,10} = [A_{2,10}^2 + B_{2,10}^2]^{1/2}$

6 Ostwald colours (o)


of maximum (m) $C_{AB,10}$ in

linear colour space ($C_{AB,2,10}, Y_{10}$)

Illumin. P00, $Y_{W,10} = 100, Y_{N,10} = 10$

Name	Range	$X_{d,10}$	$Y_{d,10}$	$Z_{d,10}$	$x_{d,10}$	$y_{d,10}$	λ_d	λ_c
R_d	567_775	73.2	48.59	8.2	0.563	0.3737	597	484
Y_d	489_775	90.23	93.93	12.77	0.4581	0.4769	571	461
G_d	489_567	27.36	55.43	12.77	0.2863	0.5799	533	533c
C_d	380_567	39.51	61.51	81.26	0.2167	0.3374	484	597
B_d	380_489	22.48	16.17	76.68	0.1949	0.1402	461	571
M_d	567_489	85.35	54.67	76.68	0.3938	0.2522	533c	533
W_d	380_775	102.3799	99.99	81.25	0.3609	0.3525	100%	
N_d	380_775	10.23	9.99	8.12	0.3609	0.3525	10%	
Z_d	380_775	18.42	18.0	14.62	0.3609	0.3525	18%	

$L^*_{10} = 60 \log[f(Y_{10,an})]$ $Y_{10,an} = [Y_{10} - 50] / 50$



$f(Y_{10,an}) = \pm [1 + 10 |Y_{10,an}|^n]$

n increases to 1 for:

1. decreasing of the contrast C
2. adjacent compared to separate colours.

Parameter:

Y_{10} & Name

Illuminant P00

$Y_{W,10} = 100, Y_{N,10} = 10$