


$XYZ_{W,10} = 97.28, 99.99, 116.14$
 $A_{2,10} = 2,5 (a_{2,10} - a_{2,n,10}) Y_{10}$
 $B_{2,10} = 2,5 B_c (b_{2,10} - b_{2,n,10}) Y_{10}$
 $a_{2,10} = a_{20} [(x_{10} - x_c) / y_{10}]$
 $b_{2,10} = b_{20} [z_{10} / y_{10}]$
 $a_{20} = 1, b_{20} = -0,4$
 $x_c = 0,110, B_c = 0,700$
 $C_{AB,2,10} = [A_{2,10}^2 + B_{2,10}^2]^{1/2}$

6 Ostwald colours (o)
 of maximum (m) $C_{AB,10}$ in
 linear colour space ($C_{AB,2,10}, Y_{10}$)
 Illumin. C00, $Y_{W,10} = 100, Y_{N,10} = 50$

Name	Range	$X_{d,10}$	$Y_{d,10}$	$Z_{d,10}$	$x_{d,10}$	$y_{d,10}$	λ_d	λ_c
R _d	561_775	79.25	70.14	58.18	0.3817	0.3379	593	481
Y _d	486_775	87.59	95.32	61.25	0.3587	0.3903	567	461
G _d	486_561	57.08	75.28	61.25	0.2948	0.3888	530	530c
C _d	380_561	66.82	80.0	116.2	0.254	0.3041	481	593
B _d	380_486	58.48	54.82	113.13	0.2582	0.2421	461	567
M _d	561_486	88.99	74.86	113.13	0.3212	0.2702	530c	530
W _d	380_775	97.28	99.99	116.14	0.3103	0.319	100%	
N _d	380_775	48.64	49.99	58.07	0.3103	0.319	50%	
Z _d	380_775	17.51	18.0	20.9	0.3103	0.319	18%	

$L^*_{10} = 60 \log[f(Y_{10,an})]$
 $Y_{10,an} = [Y_{10} - 50] / 50$



$f(Y_{10,an}) = \pm [1 + 10 |Y_{10,an}|^n]$
n increases to 1 for:
 1. decreasing of the contrast C
 2. adjacent compared to separate colours.

Parameter:
 Y_{10} & Name
 Illuminant C00
 $Y_{W,10} = 100, Y_{N,10} = 50$