

$$H^*_- = B50R_-$$

Data for maksimalfarge (Ma):

ORS20a; adapterte (a) CIELAB data
 $H^*_a \quad L^*=L^*_a \quad a^*_a \quad b^*_a \quad C^*_{ab,a} \quad h^*_{ab,a}$


HIC*_{-Ma}: B50R_100_100-

rabic* Mo:


1.0 0.0 1.0

1.0 0.0 1.0 1.0 1.0

trekantslyshet T^*



navn	$L^* = L^*_a a^*_a$	b^*_a	$C^*_{ab,a}$	$h^*_{ab,a}$	
R ₋ Ma	47.9	65.3	50.5	82.6	37
Y ₋ Ma	90.3	-10.2	91.7	92.3	96
G ₋ Ma	50.9	-62.8	34.9	71.9	150
C ₋ Ma	58.6	-30.3	-45.0	54.2	236
B ₋ Ma	25.7	31.0	-44.4	54.2	305
M ₋ Ma	48.1	75.2	-8.3	75.7	353
N ₋ Ma	18.0	0.0	0.0	0.0	0
W ₋ Ma	95.4	0.0	0.0	0.0	0
R ₋ CIE	39.9	58.7	27.9	65.0	25
Y ₋ CIE	81.2	-2.8	71.5	71.6	92
G ₋ CIE	52.2	-42.4	13.6	44.5	162
B ₋ CIE	30.5	1.4	-46.4	46.4	271



%Omfang
u*_{rel} = 92
%Regularität
g*_{H,rel} = 57
g*_{C,rel} = 58

