

$XYZ_w=95.0443, 100.0, 108.89$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a = a_2 [x/y]^{1/3}$

$b = b_2 [z/y]^{1/3}$

$a_2 = [1/X_n]^{1/3} = 0.2191$

$b_2 = -[1/Z_n]^{1/3} = -0.08376$

$n = D65$

CIELAB 76

Name and spectral range

R_m 565_770 Y_m 495_770


G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Optimal colours (o), $Y_w=100$

4 of maximum (m) C_{AB} for D65

in CIELAB diagram (a^*, b^*)



$XYZ_w=96.4228, 100.0, 82.49$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a = a_2 [x/y]^{1/3}$

$b = b_2 [z/y]^{1/3}$

$a_2 = [1/X_n]^{1/3} = 0.218$

$b_2 = -[1/Z_n]^{1/3} = -0.09188$

$n = D50$

CIELAB 76

Name and spectral range

R_m 565_770 Y_m 495_770


G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Optimal colours (o), $Y_w=100$

4 of maximum (m) C_{AB} for D50

in CIELAB diagram (a^*, b^*)



$XYZ_w = 100.932, 100.0, 64.68$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a = a_2 [x/y]^{1/3}$

$b = b_2 [z/y]^{1/3}$

$a_2 = [1/X_n]^{1/3} = 0.2147$

$b_2 = -[1/Z_n]^{1/3} = -0.09964$

$n = P40$

CIELAB 76

Name and spectral range

R_m 565_770 Y_m 495_770


G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Optimal colours (o), $Y_w = 100$

4 of maximum (m) C_{AB} for P40

in CIELAB diagram (a^* , b^*)



$XYZ_w = 109.849, 100.0, 35.58$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a = a_2 [x/y]^{1/3}$

$b = b_2 [z/y]^{1/3}$

$a_2 = [1/X_n]^{1/3} = 0.2088$

$b_2 = -[1/Z_n]^{1/3} = -0.12161$

$n = A00$

CIELAB 76

Name and spectral range

R_m 565_770 Y_m 495_770


G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Optimal colours (o), $Y_w = 100$

4 of maximum (m) C_{AB} for A00

in CIELAB diagram (a^*, b^*)



$XYZ_w=100.001, 100.0, 100.0$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a = a_2 [x/y]^{1/3}$

$b = b_2 [z/y]^{1/3}$

$a_2 = [1/X_n]^{1/3} = 0.2154$

$b_2 = -[1/Z_n]^{1/3} = -0.08617$

$n = E00$

CIELAB 76

Name and spectral range

R_m 565_770 Y_m 495_770


G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Optimal colours (o), $Y_w=100$

4 of maximum (m) C_{AB} for E00

in CIELAB diagram (a^*, b^*)



$XYZ_w=98.0718, 100.0, 118.22$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a = a_2 [x/y]^{1/3}$

$b = b_2 [z/y]^{1/3}$

$a_2 = [1/X_n]^{1/3} = 0.2168$

$b_2 = -[1/Z_n]^{1/3} = -0.08149$

$n = C00$

CIELAB 76

Name and spectral range

R_m 565_770 Y_m 495_770


G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Optimal colours (o), $Y_w=100$

4 of maximum (m) C_{AB} for C00

in CIELAB diagram (a^*, b^*)



$XYZ_w = 102.067, 100.0, 81.06$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a = a_2 [x/y]^{1/3}$

$b = b_2 [z/y]^{1/3}$

$a_2 = [1/X_n]^{1/3} = 0.2139$

$b_2 = -[1/Z_n]^{1/3} = -0.09242$

$n = P00$

CIELAB 76

Name and spectral range

R_m 565_770 Y_m 495_770


G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Optimal colours (o), $Y_w = 100$

4 of maximum (m) C_{AB} for P00

in CIELAB diagram (a^*, b^*)



$XYZ_w=97.9332, 100.0, 118.95$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a = a_2 [x/y]^{1/3}$

$b = b_2 [z/y]^{1/3}$

$a_2 = [1/X_n]^{1/3} = 0.2169$

$b_2 = -[1/Z_n]^{1/3} = -0.08133$

$n = Q00$

CIELAB 76

Name and spectral range

R_m 565_770 Y_m 495_770


G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Optimal colours (o), $Y_w=100$

4 of maximum (m) C_{AB} for Q00

in CIELAB diagram (a^*, b^*)



$XYZ_w=95.0443, 100.0, 108.89$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a = a_2 [x/y]^{1/3}$

$b = b_2 [z/y]^{1/3}$

$a_2 = [1/X_n]^{1/3} = 0.2191$

$b_2 = -[1/Z_n]^{1/3} = -0.08376$

$n = D65$

CIELAB 76

Name and spectral range

R_m 565_770 Y_m 495_770


G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Optimal colours (o), $Y_w=100$

4 of maximum (m) C_{AB} for D65

in CIELAB diagram (a^*, b^*)



$XYZ_w=96.4228, 100.0, 82.49$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a = a_2 [x/y]^{1/3}$

$b = b_2 [z/y]^{1/3}$

$a_2 = [1/X_n]^{1/3} = 0.218$

$b_2 = -[1/Z_n]^{1/3} = -0.09188$

$n = D50$

CIELAB 76

Name and spectral range

R_m 565_770 Y_m 495_770


G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Optimal colours (o), $Y_w=100$

4 of maximum (m) C_{AB} for D50

in CIELAB diagram (a^*, b^*)



$XYZ_w = 100.932, 100.0, 64.68$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a = a_2 [x/y]^{1/3}$

$b = b_2 [z/y]^{1/3}$

$a_2 = [1/X_n]^{1/3} = 0.2147$

$b_2 = -[1/Z_n]^{1/3} = -0.09964$

$n = P40$

CIELAB 76

Name and spectral range

R_m 565_770 Y_m 495_770


G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Optimal colours (o), $Y_w = 100$

4 of maximum (m) C_{AB} for P40

in CIELAB diagram (a^*, b^*)



$XYZ_w = 109.849, 100.0, 35.58$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a = a_2 [x/y]^{1/3}$

$b = b_2 [z/y]^{1/3}$

$a_2 = [1/X_n]^{1/3} = 0.2088$

$b_2 = -[1/Z_n]^{1/3} = -0.12161$

$n = A00$

CIELAB 76

Name and spectral range

R_m 565_770 Y_m 495_770


G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Optimal colours (o), $Y_w = 100$

4 of maximum (m) C_{AB} for A00

in CIELAB diagram (a^*, b^*)



$XYZ_w = 100.001, 100.0, 100.0$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a = a_2 [x/y]^{1/3}$

$b = b_2 [z/y]^{1/3}$

$a_2 = [1/X_n]^{1/3} = 0.2154$

$b_2 = -[1/Z_n]^{1/3} = -0.08617$

$n = E00$

CIELAB 76

Name and spectral range

R_m 565_770 Y_m 495_770


G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Optimal colours (o), $Y_w = 100$

4 of maximum (m) C_{AB} for E00

in CIELAB diagram (a^*, b^*)



$XYZ_w=98.0718, 100.0, 118.22$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a = a_2 [x/y]^{1/3}$

$b = b_2 [z/y]^{1/3}$

$a_2 = [1/X_n]^{1/3} = 0.2168$

$b_2 = -[1/Z_n]^{1/3} = -0.08149$

$n = C00$

CIELAB 76

Name and spectral range

R_m 565_770 Y_m 495_770


G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Optimal colours (o), $Y_w=100$

4 of maximum (m) C_{AB} for C00

in CIELAB diagram (a^*, b^*)



$XYZ_w = 102.067, 100.0, 81.06$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a = a_2 [x/y]^{1/3}$

$b = b_2 [z/y]^{1/3}$

$a_2 = [1/X_n]^{1/3} = 0.2139$

$b_2 = -[1/Z_n]^{1/3} = -0.09242$

$n = P00$

CIELAB 76

Name and spectral range

R_m 565_770 Y_m 495_770


G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Optimal colours (o), $Y_w = 100$

4 of maximum (m) C_{AB} for P00

in CIELAB diagram (a^*, b^*)



$XYZ_w=97.9332, 100.0, 118.95$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a = a_2 [x/y]^{1/3}$

$b = b_2 [z/y]^{1/3}$

$a_2 = [1/X_n]^{1/3} = 0.2169$

$b_2 = -[1/Z_n]^{1/3} = -0.08133$

$n = Q00$

CIELAB 76

Name and spectral range

R_m 565_770 Y_m 495_770

G_o 495_565 C_m 380_565

B_m 380_495 M_o 565_495

Optimal colours (o), $Y_w=100$

4 of maximum (m) C_{AB} for Q00

in CIELAB diagram (a^*, b^*)

