

$XYZ_w=95.0443, 100.0, 108.89$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = D65$

CIELAB D65

Nome e la gamma spettrale

R_m 561_770 Y_m 520_770


G_m 475_573 C_m 380_561

B_m 380_520 M_m 573_475

Ostwald colori ottimi (o), $Y_w=100$

6 di massimo (m) C_{AB} per D65

nel diagramma di chroma (a^*_{D65}, b^*_{D65})



$XYZ_w=96.4228, 100.0, 82.49$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = D50$

CIELAB D65

Nome e la gamma spettrale

R_m 561_770 Y_m 520_770


G_m 475_573 C_m 380_561

B_m 380_520 M_m 573_475

Ostwald colori ottimi (o), $Y_w=100$

6 di massimo (m) C_{AB} per D50

nel diagramma di chroma (a^*_{D65}, b^*_{D65})



$XYZ_w=100.932, 100.0, 64.68$

b^*_{D65}

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$


$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = P40$



$XYZ_w = 109.849, 100.0, 35.58$

b^*_{D65}

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.0811$

$n = A00$

CIELAB D65

Nome e la gamma spettrale

R_m 561_770 Y_m 520_770


G_m 475_573 C_m 380_561

B_m 380_520 M_m 573_475

Ostwald colori ottimi (o), $Y_w = 100$

6 di massimo (m) C_{AB} per A00

nel diagramma di chroma (a^*_{D65}, b^*_{D65})



$XYZ_w = 100.001, 100.0, 100.0$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = E00$

CIELAB D65

Nome e la gamma spettrale

R_m 561_770 Y_m 520_770


G_m 475_573 C_m 380_561

B_m 380_520 M_m 573_475

Ostwald colori ottimi (o), $Y_w = 100$

6 di massimo (m) C_{AB} per E00

nel diagramma di chroma (a^*_{D65}, b^*_{D65})



$XYZ_w=98.0718, 100.0, 118.22$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$


$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = C00$



CIE LAB D65

Nome e la gamma spettrale

R_m 561_770 Y_m 520_770

G_m 475_573 C_m 380_561

B_m 380_520 M_m 573_475

Ostwald colori ottimi (o), $Y_w=100$

6 di massimo (m) C_{AB} per C00

nel diagramma di chroma (a^*_{D65}, b^*_{D65})

$XYZ_w = 102.067, 100.0, 81.06$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = P00$

CIELAB D65

Nome e la gamma spettrale

R_m 561_770 Y_m 520_770


G_m 475_573 C_m 380_561

B_m 380_520 M_m 573_475

Ostwald colori ottimi (o), $Y_w = 100$

6 di massimo (m) C_{AB} per P00

nel diagramma di chroma (a^*_{D65}, b^*_{D65})



$XYZ_w=97.9332, 100.0, 118.95$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = Q00$

CIELAB D65

Nome e la gamma spettrale

R_m 561_770 Y_m 520_770


G_m 475_573 C_m 380_561

B_m 380_520 M_m 573_475

Ostwald colori ottimi (o), $Y_w=100$

6 di massimo (m) C_{AB} per Q00

nel diagramma di chroma (a^*_{D65}, b^*_{D65})



$XYZ_w=94.8136, 100.0, 107.33$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = D65$

CIELAB D65

Nome e la gamma spettrale

R_m 561_770 Y_m 520_770


G_m 475_573 C_m 380_561

B_m 380_520 M_m 573_475

Ostwald colori ottimi (o), $Y_w=100$

6 di massimo (m) C_{AB} per D65

nel diagramma di chroma (a^*_{D65}, b^*_{D65})



$XYZ_w=96.7256, 100.0, 81.41$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = D50$

CIELAB D65

Nome e la gamma spettrale

R_m 561_770 Y_m 520_770


G_m 475_573 C_m 380_561

B_m 380_520 M_m 573_475

Ostwald colori ottimi (o), $Y_w=100$

6 di massimo (m) C_{AB} per D50

nel diagramma di chroma (a^*_{D65}, b^*_{D65})



$XYZ_w = 101.751, 100.0, 64.44$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = P40$

CIELAB D65

Nome e la gamma spettrale

R_m 561_770 Y_m 520_770


G_m 475_573 C_m 380_561

B_m 380_520 M_m 573_475

Ostwald colori ottimi (o), $Y_w = 100$

6 di massimo (m) C_{AB} per P40

nel diagramma di chroma (a^*_{D65}, b^*_{D65})



$XYZ_w=111.15, 100.0, 35.19$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = A00$

CIELAB D65

Nome e la gamma spettrale

R_m 561_770 Y_m 520_770


G_m 475_573 C_m 380_561

B_m 380_520 M_m 573_475

Ostwald colori ottimi (o), $Y_w=100$

6 di massimo (m) C_{AB} per A00

nel diagramma di chroma (a^*_{D65}, b^*_{D65})



$XYZ_w=99.9908, 99.9999, 100.0$ b^*_{D65}

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = E00$

CIELAB D65

Nome e la gamma spettrale

R_m 561_770 Y_m 520_770


G_m 475_573 C_m 380_561

B_m 380_520 M_m 573_475

Ostwald colori ottimi (o), $Y_w=100$

6 di massimo (m) C_{AB} per E00

nel diagramma di chroma (a^*_{D65}, b^*_{D65})



$XYZ_w=97.2866, 100.0, 116.14$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = C00$

CIELAB D65

Nome e la gamma spettrale

R_m 561_770 Y_m 520_770


G_m 475_573 C_m 380_561

B_m 380_520 M_m 573_475

Ostwald colori ottimi (o), $Y_w=100$

6 di massimo (m) C_{AB} per C00

nel diagramma di chroma (a^*_{D65}, b^*_{D65})



$XYZ_w = 102.375, 100.0, 81.25$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = P00$

CIELAB D65

Nome e la gamma spettrale

R_m 561_770 Y_m 520_770


G_m 475_573 C_m 380_561

B_m 380_520 M_m 573_475

Ostwald colori ottimi (o), $Y_w = 100$

6 di massimo (m) C_{AB} per P00

nel diagramma di chroma (a^*_{D65}, b^*_{D65})



$XYZ_w=97.65, 100.0, 118.42$

$a^* = 500 (a' - a'_n) Y^{1/3}$

$b^* = 500 (b' - b'_n) Y^{1/3}$

$a' = a_2 [x/y]^{1/3}$

$b' = b_2 [z/y]^{1/3}$

$a_2 = [1/X_{D65}]^{1/3} = 0.2191$

$b_2 = -[1/Z_{D65}]^{1/3} = -0.08376$

$n = Q00$

CIELAB D65

Nome e la gamma spettrale

R_m 561_770 Y_m 520_770

G_m 475_573 C_m 380_561

B_m 380_520 M_m 573_475

Ostwald colori ottimi (o), $Y_w=100$

6 di massimo (m) C_{AB} per Q00

nel diagramma di chroma (a^*_{D65}, b^*_{D65})

