

log ΔL Leuchtdichte-Differenz- $\bullet L_g=6,3\text{cd/m}^2$
 renzschwelle

AD 26&0,1s G 6,3cd/m²; hyp2

$$\Delta L = A_1 \cdot A_2 \cdot A_3 \cdot L^t / (L^t + A_2)^2$$

$$A_1=174.4 \quad A_1=67.58$$


$$A_2=4.14 \quad A_2=3.41$$

$$A_3=0.42=t \quad A_3=0.69=t$$

$$A_6=307.2 \quad A_6=159.7$$

$$A_7=28.29 \quad A_7=5.9$$

$$\Delta=3.872 \quad \Delta=3.867$$



$\log(L/\Delta L)$ Leuchtdichte-Kontrast-
Empfindlichkeitsschwelle $L_g = 6,3 \text{ cd/m}^2$

AD 26&0,1s G 6,3cd/m²; hyp2

$$\log(L/\Delta L) = A_1 \cdot A_2 \cdot t \cdot L / (L^t + A_2)^2$$

$$A_1 = 174.4 \quad A_1 = 67.58$$


$$A_2 = 4.14 \quad A_2 = 3.41$$

$$A_3 = 0.42 = t \quad A_3 = 0.69 = t$$

$$A_6 = 207.2 \quad A_6 = 159.7$$

$$A_7 = 28.29 \quad A_7 = 5.9$$

$$\Delta = 3.772 \quad \Delta = 1.367$$



$L/\Delta L$ Leuchtdichte-Kontrast- $\bullet L_g = 6,3 \text{ cd/m}^2$
 Empfindlichkeitsschwelle

AD 26&0,1s G 6,3cd/m²; hyp2

$$L/\Delta L = A_1 \cdot A_2 \cdot t \cdot L / (L^t + A_2)^2$$

$$A_1 = 174.4 \quad A_1 = 67.58$$


$$A_2 = 4.14 \quad A_2 = 3.41$$

$$A_3 = 0.42 = t \quad A_3 = 0.69 = t$$

$$A_6 = 307.2 \quad A_6 = 159.7$$

$$A_7 = 28.29 \quad A_7 = 5.9$$

$$\Delta = 3.772 \quad \Delta = 1.367$$



T^* Leuchtdichte-Differenz-
renzschwellensumme

$AD\ 26\&0,1s\ G\ 6,3cd/m^2; hyp2$

$$T^* = A_1 \cdot L^t / (L^t + A_2)$$

$$A_1 = 174.4 \quad A_1 = 67.58$$

$$A_2 = 4.14 \quad A_2 = 3.41$$

$$A_3 = 0.42 = t \quad A_3 = 0.69 = t$$

$$A_6 = 307.2 \quad A_6 = 159.7$$

$$A_7 = 28.29 \quad A_7 = 5.9$$

$$\Delta = 3.772 \quad \Delta = 1.367$$

$$L_g = 6,3cd/m^2$$

