

log ΔL Leuchtdichte-Differenz- $\bullet L_g=63\text{cd/m}^2$
 renzschwelle

AD 26s G 63cd/m²; hyp2

$$\Delta L = A_1 \cdot A_2 \cdot A_3 \cdot L^t / (L^t + A_2)^2$$

$$A_1=168.3$$


$$A_2=17.36$$

$$A_3=0.8=t$$

$$A_6=2337.0$$

$$A_7=35.41$$

$$\Delta=4.936$$



$\log(L/\Delta L)$ Leuchtdichte-Kontrast
Empfindlichkeitsschwelle

AD 26s G 63cd/m²; hyp2

$$\log(L/\Delta L) = A_1 \cdot A_2 \cdot t \cdot L^{A_3} \cdot (E + A_2)^2$$

$$A_1 = 168.3$$


$$A_2 = 17.36$$

$$A_3 = 0.8 = t$$

$$A_6 = 2337.0$$

$$A_7 = 35.41$$

$$\Delta = 4.936$$



$L/\Delta L$ Leuchtdichte-Kontrast- $\bullet L_g = 63 \text{cd/m}^2$
 Empfindlichkeitsschwelle

$AD\ 26s\ G\ 63 \text{cd/m}^2; \text{hyp2}$

$$L/\Delta L = A_1 \cdot A_2 \cdot t \cdot L / (L^t + A_2)^2$$

$$A_1 = 168.3$$


$$A_2 = 17.36$$

$$A_3 = 0.8 = t$$

$$A_6 = 2337.0$$

$$A_7 = 35.41$$

$$\Delta = 4.936$$



T^* Leuchtdichte-Differenz-
renzschwellensumme

$AD\ 26s\ G\ 63cd/m^2; hyp2$

$$T^* = A_1 \cdot L^t / (L^t + A_2)$$

$$A_1 = 168.3$$

$$A_2 = 17.36$$

$$A_3 = 0.8 = t$$

$$A_6 = 2337.0$$

$$A_7 = 35.41$$

$$\Delta = 4.936$$

$$L_g = 63cd/m^2$$

