

log ΔL Leuchtdichte-Differenz-
renzschwelle • $L_g = 6,3 \text{ cd/m}^2$

04 26&0,1s A 6,3cd/m²; hyp3

$$\Delta L = A_1 \cdot A_2 \cdot A_3 \cdot L^t / (L^t + A_2)^2$$

$$A_1 = 201.9 \quad A_1 = 81.89$$


$$A_2 = 12.88 \quad A_2 = 5.59$$

$$A_3 = 0.8=t \quad A_3 = 0.8=t$$

$$A_6 = 2080.0 \quad A_6 = 366.6$$

$$A_7 = 24.4 \quad A_7 = 8.6$$

$$\Delta = 91.01 \quad \Delta = 7.289$$



$L/\Delta L$ Leuchtdichte-Kontrast- $\bullet L_g=6,3\text{cd/m}^2$
 Empfindlichkeitsschwelle

04 26&0,1s A 6,3cd/m²; hyp3

$$L/\Delta L = A_1 \cdot A_2 \cdot t \cdot L / (L^t + A_2)^2$$

$$A_1=201.9 \quad A_1=81.89$$


$$A_2=12.88 \quad A_2=5.59$$

$$A_3=0.8=t \quad A_3=0.8=t$$

$$A_6=2080.0 \quad A_6=366.6$$

$$A_7=24.4 \quad A_7=8.6$$

$$\Delta=91.01 \quad \Delta=7.289$$



T^* Leuchtdichte-Differenz- renzschwellensumme

• $L_g = 6,3 \text{ cd/m}^2$

80 04 26 & 0,1 s A 6,3 cd/m²; hyp3

$$T^* = A_1 \cdot L^t / (L^t + A_2)$$

$$A_1 = 201.9 \quad A_1 = 81.89$$

$$A_2 = 12.88 \quad A_2 = 5.59$$

$$A_3 = 0.8 = t \quad A_3 = 0.8 = t$$

$$A_6 = 2080.0 \quad A_6 = 366.6$$

$$A_7 = 24.4 \quad A_7 = 8.6$$

$$\Delta = 91.01 \quad \Delta = 7.285$$

