

Munsell (Renotation)-Buntheit  $C = 2$  und Helligkeit (Value)  $V = 1, 5$  und  $9$   
 in Farbardiagramm ( $a_2(F,M)$ ,  $b_2(F,M)$ )


$$b_2(F,M) = b_2(F) - b_2(M)$$

$$a_{20} = 1.0, b_{20} = -0.4$$

$$x_c = 0.11, n = 0.341, B_c = 1.0$$

$$a_2 = a_{20} (x - x_c) / y$$

$$b_2 = b_{20} B_c z / y$$



$$a_2(F,M) = a_2(F) - a_2(M)$$

| $V$ | $a_2(F,M)$ | $b_2(F,M)$ |
|-----|------------|------------|
| 1   | 0.0        | 0.0        |
| 2   | 0.0        | 0.0        |
| 3   | 0.0        | 0.0        |
| 4   | 0.0        | 0.0        |
| 5   | 0.0        | 0.0        |
| 6   | 0.0        | 0.0        |
| 7   | 0.0        | 0.0        |
| 8   | 0.0        | 0.0        |
| 9   | 0.0        | 0.0        |
| M   | 0.0        | 0.0        |

$$x_M = 0.313$$

$$y_M = 0.324$$

$$V_M = 25.30$$