


Munsell (Renotation)-Buntheit C = 2 und Helligkeit (Value) V = 1, 5 und 9
in Farbartdiagramm ($p_2(F,M)$, $q_2(F,M)$) $p_{20}=2.806$, $q_{20}=1.0$

$$q_2(F,M) = q_2(F) - q_2(M) \quad x_c=0.11, B_c=1.0$$

$$p_2 = p_{20} B_c [3,0757(x-x_c) - 2,5702 y - 0,0960] / y$$

$$q_2 = q_{20} [1,9907(x-x_c) + 3,8617 y - 2,4046] / y$$




cgd90-5a

Munsell (Renotation)-Buntheit C = 2 und Helligkeit (Value) V = 1, 5 und 9
in Buntheitsdiagramm ($p_2^*(F,M)$, $q_2^*(F,M)$) $p_{20}=2.806$, $q_{20}=1.0$

$$q_2^*(F,M) = c_Y [q_2(F) - (M)]] = c_Y q_2(F,M) \quad x_c=0.11, B_c=1.0, c_Y=0,91 Y^{0,341}$$

$$p_2 = p_{20} B_c [3,0757(x-x_c) - 2,5702 y - 0,0960] / y$$

$$q_2 = q_{20} [1,9907(x-x_c) + 3,8617 y - 2,4046] / y$$




cgd90-6a

Munsell (Renotation)-Buntheit C = 2 und Helligkeit (Value) V = 1, 5 und 9
in Farbartdiagramm ($p_2(F,M)$, $q_2(F,M)$) $p_{20}=2.806$, $q_{20}=1.0$

$$q_2(F,M) = q_2(F) - q_2(M) \quad x_c=0.11, B_c=0.8$$

$$p_2 = p_{20} B_c [3,0757(x-x_c) - 2,5702 y - 0,0960] / y$$

$$q_2 = q_{20} [1,9907(x-x_c) + 3,8617 y - 2,4046] / y$$




cgd90-7a

Munsell (Renotation)-Buntheit C = 2 und Helligkeit (Value) V = 1, 5 und 9
in Buntheitsdiagramm ($p_2^*(F,M)$, $q_2^*(F,M)$) $p_{20}=2.806$, $q_{20}=1.0$

$$q_2^*(F,M) = c_Y [q_2(F) - (M)]] = c_Y q_2(F,M) \quad x_c=0.11, B_c=0.8, c_Y=0,91 Y^{0,341}$$

$$p_2 = p_{20} B_c [3,0757(x-x_c) - 2,5702 y - 0,0960] / y$$

$$q_2 = q_{20} [1,9907(x-x_c) + 3,8617 y - 2,4046] / y$$



cgd90-8a