

Optimalfarben (o) RYGCBM von maximalem (m) $C_{AB,10}$; D65, $Y_m=510_770$, LABHNU1_79

<i>Code</i> D65	L^*_{10}	A^*_{10}	B^*_{10}	$C^*_{ab,10}$	a'_{10}	b'_{10}	$h_{ab,10}$	i_d	λ^*_d	i_c	λ^*_c
R _{me} 570_770	62.44	70.17	66.75	96.85	0.1745	-0.0469	43.5	38	592	14	473
Y _{me} 510_770	88.89	-1.92	87.39	87.41	0.1289	-0.0478	91.2	33	565	13	465
G _{me} 470_570	79.44	-72.31	42.58	83.92	0.0919	-0.0672	149.5	21	508	8	440
C _m 380_570	81.53	-45.38	-22.81	50.79	0.1066	-0.1012	206.6	15	477	44	622
B _{me} 380_510	47.53	5.28	-78.32	78.5	0.134	-0.1511	273.8	13	465	32	564
M _m 570_470	65.48	99.23	-35.65	105.44	0.1907	-0.1114	340.2	2	413	19	498
R _o 570_440	63.05	86.25	6.33	86.49	0.1844	-0.0855	4.2	-1	484c	16	484
G _o 520_570	71.8	-58.74	70.82	92.02	0.0964	-0.0491	129.6	27	536	11	455
W ₁ 380_770	96.0	0.0	0.0	0.0	0.1298	-0.0895	0.0	-1	486c	17	486