

A choose of a value different "0.50" changes the grey sample and surround. Beginners often have difficulties to choose on an appropriate value. Therefore it is recommended for beginners to proceed with image 2. After a restart of the experiment, a value different "0.50" may be used.

adjust visual equal difference for one of 3 steps

Output (9 steps) adjusted spacing $0 < r_{gb}^{in} < 1$

go to next image 2

one experimental value: e_{58}

equally spaced $0 < r_{gb}^{in} < 1$ Input (9 steps)

heq80-1a, image 1, produce equal visual difference between Magenta M – Magenta Mw – White

9 step series based only on the visual adjustment of image 1 with value "0.50" or different

adjust visual equal difference for two of 5 steps

Output (9 steps) adjusted spacing $0 < r_{gb}^{in} < 1$

go to next image 3

two experimental values: e_{64}, e_{48}

equally spaced $0 < r_{gb}^{in} < 1$ Input (9 steps)

heq80-2a, image 2, produce equal visual difference between two of five steps

9 step series based only on the visual adjustment of image 1 with value "0.50" or different

adjust visual equal difference for four of 9 steps

Output (9 steps) adjusted spacing $0 < r_{gb}^{in} < 1$

go to next image 4

four experimental values: $e_{62}, e_{24}, e_{46}, e_{68}$

save 7 data above as text

equally spaced $0 < r_{gb}^{in} < 1$ Input (9 steps)

heq80-3a, image 3, produce equal visual difference between four of nine steps

heq80-3n

9 step series based only on the visual adjustment of image 1 with value "0.50" or different

9 step series based on all visual adjustments used for output linearization

calculation with visual experimental (c) data adjusted above

save 7 data above as text

save 9 data below as text

grey example difference visible? $0.25 + 0.06$ adjust threshold $0.25 + 0.00$ no change

adjust and proof threshold of the linearized output

restart with image 1

heq80-4a, image 4, adjust visual threshold (+0.047) of 9 steps; all equal?

A choose of a value different "0.50" changes the grey sample and surround. Beginners often have difficulties to choose on an appropriate value. Therefore it is recommended for beginners to proceed with image 2. After a restart of the experiment, a value different "0.50" may be used.

adjust visual equal difference for one of 3 steps

Output (9 steps) adjusted spacing $0 < r_{gb}^{in} < 1$

go to next image 2

one experimental value: e_{58}

equally spaced $0 < r_{gb}^{in} < 1$ Input (9 steps)

heq81-1a, image 1, produce equal visual difference between Magenta M – Magenta Mw – White

9 step series based only on the visual adjustment of image 1 with value "0.50" or different

adjust visual equal difference for two of 5 steps

Output (9 steps) adjusted spacing $0 < r_{gb}^{in} < 1$

go to next image 3

two experimental values: e_{64}, e_{48}

equally spaced $0 < r_{gb}^{in} < 1$ Input (9 steps)

heq81-2a, image 2, produce equal visual difference between two of five steps

9 step series based only on the visual adjustment of image 1 with value "0.50" or different

adjust visual equal difference for four of 9 steps

Output (9 steps) adjusted spacing $0 < r_{gb}^{in} < 1$

go to next image 4

four experimental values: $e_{62}, e_{24}, e_{46}, e_{68}$

save 7 data above as text

equally spaced $0 < r_{gb}^{in} < 1$ Input (9 steps)

heq81-3a, image 3, produce equal visual difference between four of nine steps

heq81-3n

9 step series based only on the visual adjustment of image 1 with value "0.50" or different

9 step series based on all visual adjustments used for output linearization

calculation with visual experimental (c) data adjusted above

save 7 data above as text

save 9 data below as text

grey example difference visible? $0.25 + 0.06$ adjust threshold $0.25 + 0.00$ no change

adjust and proof threshold of the linearized output

restart with image 1

heq81-4a, image 4, adjust visual threshold (+0.047) of 9 steps; all equal?

A choose of a value different "0.50" changes the grey sample and surround. Beginners often have difficulties to choose on an appropriate value. Therefore it is recommended for beginners to proceed with image 2. After a restart of the experiment, a value different "0.50" may be used.

adjust visual equal difference for one of 3 steps

Output (9 steps) adjusted spacing $0 < r_{gb}^{in} < 1$

go to next image 2

one experimental value: e_{58}

equally spaced $0 < r_{gb}^{in} < 1$ Input (9 steps)

heq80-5a, image 1, produce equal visual difference between Magenta M – Magenta Mw – White

9 step series based only on the visual adjustment of image 1 with value "0.50" or different

adjust visual equal difference for two of 5 steps

Output (9 steps) adjusted spacing $0 < r_{gb}^{in} < 1$

go to next image 3

two experimental values: e_{64}, e_{48}

equally spaced $0 < r_{gb}^{in} < 1$ Input (9 steps)

heq80-6a, image 2, produce equal visual difference between two of five steps

A choose of a value different "0.50" changes the grey sample and surround. Beginners often have difficulties to choose on an appropriate value. Therefore it is recommended for beginners to proceed with image 2. After a restart of the experiment, a value different "0.50" may be used.

adjust visual equal difference for one of 3 steps

Output (9 steps) adjusted spacing $0 < r_{gb}^{in} < 1$

go to next image 2

one experimental value: e_{58}

equally spaced $0 < r_{gb}^{in} < 1$ Input (9 steps)

heq81-5a, image 1, produce equal visual difference between Magenta M – Magenta Mw – White

9 step series based only on the visual adjustment of image 1 with value "0.50" or different

adjust visual equal difference for two of 5 steps

Output (9 steps) adjusted spacing $0 < r_{gb}^{in} < 1$

go to next image 3

two experimental values: e_{64}, e_{48}

equally spaced $0 < r_{gb}^{in} < 1$ Input (9 steps)

heq81-6a, image 2, produce equal visual difference between two of five steps

9 step series based only on the visual adjustment of image 1 with value "0.50" or different

adjust visual equal difference for four of 9 steps

Output (9 steps) adjusted spacing $0 < r_{gb}^{in} < 1$

go to next image 4

four experimental values: $e_{62}, e_{24}, e_{46}, e_{68}$

save 7 data above as text

equally spaced $0 < r_{gb}^{in} < 1$ Input (9 steps)

heq80-7a, image 3, produce equal visual difference between four of nine steps

heq80-7n

9 step series based only on the visual adjustment of image 1 with value "0.50" or different

9 step series based on all visual adjustments used for output linearization

calculation with visual experimental (c) data adjusted above

save 7 data above as text

save 9 data below as text

grey example difference visible? $0.25 + 0.06$ adjust threshold $0.25 + 0.00$ no change

adjust and proof threshold of the linearized output

restart with image 1

heq80-8a, image 4, adjust visual threshold (+0.047) of 9 steps; all equal?

9 step series based only on the visual adjustment of image 1 with value "0.50" or different

adjust visual equal difference for four of 9 steps

Output (9 steps) adjusted spacing $0 < r_{gb}^{in} < 1$

go to next image 4

four experimental values: $e_{62}, e_{24}, e_{46}, e_{68}$

save 7 data above as text

equally spaced $0 < r_{gb}^{in} < 1$ Input (9 steps)

heq81-7a, image 3, produce equal visual difference between four of nine steps

heq81-7n

9 step series based only on the visual adjustment of image 1 with value "0.50" or different

9 step series based on all visual adjustments used for output linearization

calculation with visual experimental (c) data adjusted above

save 7 data above as text

save 9 data below as text

grey example difference visible? $0.25 + 0.06$ adjust threshold $0.25 + 0.00$ no change

adjust and proof threshold of the linearized output

restart with image 1

heq81-8a, image 4, adjust visual threshold (+0.047) of 9 steps; all equal?