


$F_{ab}(x_r) = \text{unbunte Rezeptorerregungen } N, W, N+W$

$$F^*(x_r) = b \frac{e^{x_r/a} - e^{-x_r/a}}{e^{x_r/a} + e^{-x_r/a}}$$

$$a_0=-1,00, b_0=1,00$$

$$a_1=1,00, b_1=1,00$$




hgb00-5a

$F_{ab}(x_r) = \text{unbunte Rezeptorerregungen } N, W, N+W$

$$F^*(x_r) = b \frac{e^{x_r/a} - e^{-x_r/a}}{e^{x_r/a} + e^{-x_r/a}}$$

$$a_0=-0,71, b_0=1,00$$

$$a_1=1,00, b_1=2,00$$




hgb00-7n

$F_{ab}(x_r) = \text{unbunte Rezeptorerregungen } N, W, N+W$

$$F^*(x_r) = b \frac{e^{x_r/a} - e^{-x_r/a}}{e^{x_r/a} + e^{-x_r/a}}$$

$$a_0=-1,00, b_0=1,00$$

$$a_1=1,00, b_1=2,00$$




hgb00-6a

$F_{ab}(x_r) = \text{unbunte Rezeptorerregungen } N, W, N+W$

$$F^*(x_r) = b \frac{e^{x_r/a} - e^{-x_r/a}}{e^{x_r/a} + e^{-x_r/a}}$$

$$a_0=-0,71, b_0=1,00$$

$$a_1=1,00, b_1=2,00$$



hgb00-8a

hgb00-7n